


# NAVAL SUBMARINE MEDICAL RESEARCH LABORATORY

## SUBMARINE BASE, GROTON, CONN.



REPORT NUMBER 954

REFINEMENTS OF FORMULAS FOR RECEPTOR SENSITIVITY

by

James A. Worthey and David F. Neri

Naval Medical Research and Development Command  
Research Work Unit MR000.01.01-5087

Released by:

William C. Milroy, CAPT, MC, USN  
Commanding Officer  
Naval Submarine Medical Research Laboratory

31 January 1982

## SUMMARY PAGE

### PROBLEM

To improve a method for approximating functions of wavelengths which is applicable to computer calculations related to vision.

### FINDINGS

Shifting the origin of the wavelength scale to the center of the region of interest resulted in a significant reduction of roundoff errors.

### APPLICATION

Use of the method presented enables researchers to use convenient approximation methods for calculations related to vision, without restriction on the computer or calculator employed. A previously published method would fail on some machines due to roundoff error.

### ADMINISTRATIVE INFORMATION

This investigation was conducted as part of the Naval Submarine Medical Research Laboratory Work Unit MR000.01.01-5087 -- "Under-stimulation of the red-green opponent system by fluorescent light." This manuscript was submitted for review on 28 April 1981, approved for publication on 13 May 1981 and designated as NSMRL Report No. 954.

# Refinements of formulas for receptor sensitivity

James A. Worthey and David F. Neri

Naval Submarine Medical Research Laboratory, Naval Submarine Base, Groton, Connecticut 06349

Received May 19, 1981

When it is desired to fit a polynomial to a function of wavelength, roundoff errors may be minimized by shifting the origin of the independent variable (wavelength or frequency) to the center of the region of interest.

As mathematical modeling of visual functions becomes more successful, it is naturally desirable that independent workers should be able to reproduce a given model easily and accurately. Boynton and Wisowaty<sup>1</sup> have taken an important step in this direction by calculating formulas that approximate a certain set of receptor-sensitivity functions put forth by Smith and Pokorny.<sup>2</sup> In this Letter, we present some simple manipulations of the Boynton and Wisowaty results that will make them easier to use. The effect of the transformation that we make is to reduce roundoff errors, but the goal is *not* to make the answers more accurate. Rather, the use of formulas that are more immune to roundoff trouble removes the requirement for a computer that carries 10 significant digits and the need for coefficients carried out to 10 digits.

The step that we take is to shift the origin of the independent variable to the center of the region of interest. This traditional device of applied mathematics has general application when a polynomial is to be fitted to a function of wavelength. If the device were applied at the beginning of a regression calculation, it would reduce or eliminate roundoff error at all stages. Finally, in the same spirit of making calculations more workable and routine, we make explicit the relation of the present receptor sensitivities to Judd's  $\bar{x}'$ ,  $\bar{y}'$ , and  $\bar{z}'$  and to the receptor sensitivities incorporated in the model of Guth *et al.*<sup>3</sup>

In discussing the use of their formulas, Boynton and Wisowaty observe that "... for some machines rounding errors may cause problems with the fourth-order polynomials (it is essential to retain all five decimal places for the coefficients of Table I)." <sup>4</sup> By "five decimal places," Boynton and Wisowaty mean some 8–10 significant digits on most of the coefficients. Roundoff errors occur because the formulas as given depend on the subtraction of nearly equal numbers. For instance, one step in the evaluation of the green receptor function involves the polynomial evaluation

$$\log a = a_4\lambda^4 + a_3\lambda^3 + a_2\lambda^2 + a_1\lambda + a_0. \quad (1)$$

Using coefficients  $a_0$  through  $a_4$ , as provided by Boynton and Wisowaty, and  $\lambda = 0.5 \mu\text{m}$  leads to the sum

$$\begin{aligned} \log a &= -1747.361113 + 6533.32095 - 9149.3564 \\ &\quad + 5695.5237 - 1332.9406 \\ &= -0.813463. \end{aligned} \quad (2)$$

To shift the origin of wavelength,<sup>5</sup> we define the new variable

$$l = \lambda - \lambda_0. \quad (3)$$

As a practical matter, we put the new origin in the middle of the visible spectrum:

$$\lambda_0 = 0.55 \mu\text{m}. \quad (4)$$

Equation (1) may be replaced by a polynomial in  $l$ :

$$\log a = \alpha_4 l^4 + \alpha_3 l^3 + \alpha_2 l^2 + \alpha_1 l + \alpha_0. \quad (5)$$

In general, to transform the coordinates of a polynomial of degree  $N$ , the new coefficients  $\alpha_k$  can be calculated from the old  $a_k$  by

$$\alpha_k = \sum_{j=0}^{N-k} a_{j+k} \frac{(j+k)!}{j!k!} \lambda_0^j, \quad (6)$$

where  $k = 0 \dots N$ . With  $\lambda_0 = 0.55 \mu\text{m}$  and  $\lambda = 0.5 \mu\text{m}$ , the evaluation of Eq. (5) involves the sum

$$\begin{aligned} \log a &= -0.1747 + 1.1551 - 2.7525 + 2.0055 - 1.0470 \\ &= -0.8136. \end{aligned} \quad (7)$$

The answer is essentially the same, but roundoff error is less likely to occur since we are no longer adding and subtracting quantities  $10^4$  times as large as the answer. Any time one wishes to fit a polynomial to data that have a limited domain of the independent variable not centered on the origin, a similar reduction of roundoff problems can be expected from a transformation such as Eq. (3), with  $\lambda_0$  in the center of the domain.

In the case of the red- and blue-sensitivity functions, Boynton and Wisowaty found it necessary to fit polynomials in frequency to functions expressed as quantum sensitivity. In this case, the origin of frequency must be shifted. We define

$$f = \nu - \nu_0. \quad (8)$$

As a practical matter, we choose

$$\nu_0 = 5.9 \quad (9)$$

in units of  $10^{14}$  Hz—the convenient unit selected by Boynton and Wisowaty.

The working equations now become Eqs. (3), (4), (8), and (9) above, and Eqs. (10)–(22) below. Green-catching sensitivity is given by

$$G = (a^n + b^n)^{1/n}, \quad (10)$$

$$\log a = \alpha_4 l^4 + \alpha_3 l^3 + \alpha_2 l^2 + \alpha_1 l + \alpha_0, \quad (11)$$

$$\log b = \beta_4 l^4 + \beta_3 l^3 + \beta_2 l^2 + \beta_1 l + \beta_0. \quad (12)$$

**Table 1. Values of  $n$  for Eq. (10) for  $G$ , and for Eq. (14) for  $R$  and  $B$ , and the coefficients for Eqs. (11), (12), and (15)–(17)**

Exponents		
$R$	$G$	$B$
$n_1 = 2.25$	$n = 1.45$	$n_1 = 8.00$
$n_2 = 8.50$	–	$n_2 = 5.00$
Coefficients		
$R$	$G$	$B$
$\alpha_3 = -0.02467$	$\alpha_4 = -27960.$	$\alpha_3 = 0.6877$
$\alpha_2 = -0.9920$	$\alpha_3 = -9241.$	$\alpha_2 = -4.020$
$\alpha_1 = 1.432$	$\alpha_2 = -1101.$	$\alpha_1 = 5.691$
$\alpha_0 = -2.087$	$\alpha_1 = -40.11$	$\alpha_0 = -2.053$
–	$\alpha_0 = -1.047$	–
$\beta_3 = 0.5818$	$\beta_4 = -1706.$	$\beta_3 = -0.3913$
$\beta_2 = -1.042$	$\beta_3 = 1190.$	$\beta_2 = -1.305$
$\beta_1 = -1.403$	$\beta_2 = -319.7$	$\beta_1 = 2.431$
$\beta_0 = -0.5700$	$\beta_1 = 3.979$	$\beta_0 = -0.7289$
–	$\beta_0 = -0.4519$	–
$\gamma_3 = 0.6602$	–	$\gamma_3 = -11.05$
$\gamma_2 = -0.6721$	–	$\gamma_2 = -33.31$
$\gamma_1 = -1.883$	–	$\gamma_1 = -29.86$
$\gamma_0 = -0.9875$	–	$\gamma_0 = -11.16$

**Table 2. Values of  $n$  and Coefficients for Eqs. (19)–(22) To Be Used in Calculations of Eye Lens and Macular Pigment Density**

Exponents	
$n_1 = 15.00$	
$n_2 = 7.50$	
$n_3 = 4.00$	
Coefficients	
$\alpha_3 = -10130.$	$\gamma_3 = -928.0$
$\alpha_2 = -3596.$	$\gamma_2 = -485.1$
$\alpha_1 = -432.8$	$\gamma_1 = -56.18$
$\alpha_0 = -16.91$	$\gamma_0 = -1.295$
$\beta_3 = -6684.$	$\delta_3 = 14.93$
$\beta_2 = -2243.$	$\delta_2 = 2.715$
$\beta_1 = -244.5$	$\delta_1 = -1.119$
$\beta_0 = -8.016$	$\delta_0 = 0.0716$

Red-catching sensitivity  $R$  and blue-catching sensitivity  $B$  are given by

$$R \text{ or } B = Q(\lambda/555), \quad (13)$$

where  $\lambda$  is in nanometers,

$$Q = [(a^{n_1} + b^{n_1})^{n_2/n_1} + c^{n_2}]^{1/n_2}, \quad (14)$$

and

$$\log a = \alpha_3 f^3 + \alpha_2 f^2 + \alpha_1 f + \alpha_0, \quad (15)$$

$$\log b = \beta_3 f^3 + \beta_2 f^2 + \beta_1 f + \beta_0, \quad (16)$$

$$\log c = \gamma_3 f^3 + \gamma_2 f^2 + \gamma_1 f + \gamma_0. \quad (17)$$

Table 1 gives the required constants.

For the density of eye lens plus macular pigment,

$$D = \{[(a^{n_1} + b^{n_1})^{n_2/n_1} + c^{n_2}]^{n_3/n_2} + d^{n_3}\}^{1/n_3}, \quad (18)$$

where the equations for  $a$ ,  $b$ ,  $c$ , and  $d$  are

$$\log a = \alpha_3 l^3 + \alpha_2 l^2 + \alpha_1 l + \alpha_0, \quad (19)$$

$$\log b = \beta_3 l^3 + \beta_2 l^2 + \beta_1 l + \beta_0, \quad (20)$$

$$\log c = \gamma_3 l^3 + \gamma_2 l^2 + \gamma_1 l + \gamma_0, \quad (21)$$

$$\log d = \delta_3 l^3 + \delta_2 l^2 + \delta_1 l + \delta_0. \quad (22)$$

Table 2 gives the required constants.

The polynomial coefficients have been tabulated only to four significant digits. This degree of precision was chosen in a systematic way. We wrote a computer program to tabulate calculated  $R$ ,  $G$ ,  $B$ , and  $D$  with the polynomial coefficients rounded to 2, 3, 4, . . . up to 8 significant digits. Six-digit coefficients give nearly perfect agreement with the Boynton and Wisowaty calculated values; more digits give perfect agreement. Four-significant-digit precision in the coefficients gives discrepancies from those earlier calculated values of a few units in the fourth decimal place—a precision of about 1 in  $10^4$ . This should be tolerable, since Boynton and Wisowaty find their fits to the Smith and Pokorny functions to have precision of the order of 2 in  $10^2$ .

## SOME RELATED DATA

Taking the inverse of the transformation matrix derived by Smith and Pokorny<sup>2</sup> gives the expression to recover Judd's modified  $\bar{x}$ ,  $\bar{y}$ ,  $\bar{z}$ :

$$\begin{pmatrix} \bar{x}' \\ \bar{y}' \\ \bar{z}' \end{pmatrix} = \begin{pmatrix} 2.9448 & -3.5010 & 0.2118 \\ 1.0000 & 1.0000 & 0.0000 \\ 0.0000 & 0.0000 & 1.0000 \end{pmatrix} \begin{pmatrix} R \\ G \\ B \end{pmatrix}. \quad (23)$$

This provides smooth versions of the Judd functions for applications where this might be convenient.

The equations above give  $(R, G, B)$  with maxima of (0.6442, 0.3912, 1.6431) at wavelengths of (562, 542, 440) nm. Guth *et al.* utilize essentially the same  $(R, G, B)$  as a starting point for their vector model for normal and dichromatic color vision,<sup>3</sup> but with the functions adjusted to have maximum values of 1.0. To rescale  $R$ ,  $G$ , and  $B$  to have maxima of unity, multiply the expressions given above [Eqs. (10) and (13)] by 1.5523, 2.556, and 0.6086, respectively.

## REFERENCES

1. R. M. Boynton and J. J. Wisowaty, "Equations for chromatic discrimination models," J. Opt. Soc. Am. **70**, 1471–1476 (1980).
2. V. C. Smith and J. Pokorny, "Spectral sensitivity of the foveal cone photopigments between 400 and 500 nm," Vision Res. **15**, 161–171 (1975).
3. S. L. Guth, R. W. Massof, and T. Benzschawel, "Vector model for normal and dichromatic color vision," J. Opt. Soc. Am. **70**, 197–212 (1980).
4. R. M. Boynton and J. J. Wisowaty, Ref. 1, p. 1475.
5. J. H. Kindle, *Analytic Geometry* (Schaum, New York, 1950), p. 66.
6. G. W. Wyszecki and W. S. Stiles, *Color Science* (Wiley, New York, 1967), p. 436.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NSMRL Report 954	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) "Refinements of formulas for receptor sensitivity"		5. TYPE OF REPORT & PERIOD COVERED Interim report
7. AUTHOR(s) J. A. Worthey and D. F. Neri		6. PERFORMING ORG. REPORT NUMBER NSMRL Report 954
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Submarine Medical Research Laboratory Box 900 Naval Submarine Base Nlon Groton, Connecticut 06349		8. CONTRACT OR GRANT NUMBER(s)
10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS MR000.01.01-5087		11. CONTROLLING OFFICE NAME AND ADDRESS same as 9
12. REPORT DATE 31 January 1982		13. NUMBER OF PAGES 2
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Naval Medical Research and Development Command National Naval Medical Center Bethesda, Maryland 20814		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) receptor sensitivity; roundoff errors;		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  When it is desired to fit a polynomial to a function of wavelength, roundoff errors may be minimized by shifting the origin of the independent variable (wavelength or frequency) to the center of the region of interest.		

